Please provide complete and well-written solutions to the following exercises.

Due September 4, 9AM, to be submitted in brightspace.

Homework 1

Exercise 1. Estimate the probability that 1000000 coin flips of fair coins will result in more than $501,000$ heads, using the Central Limit Theorem. (Some of the following integrals may be relevant: $\int_{-\infty}^{0} e^{-t^2/2}dt/\sqrt{2\pi} = 1/2$, $\int_{-\infty}^{1} e^{-t^2/2}dt/\sqrt{2\pi} \approx .8413$, $\int_{-\infty}^{2} e^{-t^2/2}dt/\sqrt{2\pi} \approx .9772$, $\int_{-\infty}^{3} e^{-t^2/2}dt/\sqrt{2\pi} \approx .9987$.) (Hint: use Bernoulli random variables.)

Exercise 2 (Numerical Integration)**.** In computer graphics in video games, etc., various integrations are performed in order to simulate lighting effects. Here is a way to use random sampling to integrate a function in order to quickly and accurately render lighting effects. Let $\Omega = [0,1]$, and let $\mathbf{P}$ be the uniform probably law on Ω, so that if $0 \leq a < b \leq 1$, we have $\mathbf{P}([a,b]) = b - a$. Let $X_1, \ldots, X_n$ be independent random variables such that $\mathbf{P}(X_i \in [a,b]) = b - a$ for all $0 \leq a < b \leq 1$, for all $i \in \{1, \ldots, n\}$. Let $f\colon [0,1] \to \mathbf{R}$ be a continuous function we would like to integrate. Instead of integrating f directly, we instead compute the quantity

$$\frac{1}{n}\sum_{i=1}^{n} f(X_i).$$

Show that

$$\lim_{n\to\infty} \mathbf{E}\left(\frac{1}{n}\sum_{i=1}^{n} f(X_i)\right) = \int_0^1 f(t)dt.$$

$$\lim_{n\to\infty} \operatorname{var}\left(\frac{1}{n}\sum_{i=1}^{n} f(X_i)\right) = 0.$$

That is, as n becomes large, $\frac{1}{n}\sum_{i=1}^{n} f(X_i)$ is a good estimate for $\int_0^1 f(t)dt$.

Exercise 3. Let $X := (X_1, \ldots, X_n)$ be a random sample of size n from a binomial distribution with parameters n and p. Here n is a positive (known) integer and $0 < p < 1$ is unknown. (That is, $X_1, \ldots, X_n$ are i.i.d. and X_1 is a binomial random variable with parameters n and p, so that $\mathbf{P}(X_1 = k) = \binom{n}{k}p^k(1-p)^{n-k}$ for all integers $0 \leq k \leq n$.)

You can freely use that $\mathbf{E}X_1 = np$ and $\operatorname{Var}X_1 = np(1-p)$.

- Computer the Fisher information $I_X(p)$ for any $0 < p < 1$.
  (Consider n to be fixed.)
- Let Z be an unbiased estimator of p^2 (assume that Z is a function of $X_1, \ldots, X_n$). State the Cramér-Rao inequality for Z.
- Let W be an unbiased estimator of $1/p$ (assume that W is a function of $X_1, \ldots, X_n$). State the Cramér-Rao inequality for W.

Exercise 4. Let $X_1, \ldots, X_n$ be a random sample of size n from a Poisson distribution with unknown parameter $\lambda > 0$. (So, $\mathbf{P}(X_1 = k) = e^{-\lambda}\lambda^k/k!$ for all integers $k \geq 0$.)

- Find an MLE (maximum likelihood estimator) for λ.
- Is the MLE you found unique? That is, could there be more than one MLE for this problem?